

Research Article

Generalizations of Shafer-Fink-Type Inequalities for the Arc Sine Function

Wenhai Pan and Ling Zhu

Department of Mathematics, Zhejiang Gongshang University, Hangzhou, Zhejiang 310018, China

Correspondence should be addressed to Ling Zhu, zhuling0571@163.com

Received 29 December 2008; Revised 9 March 2009; Accepted 28 April 2009

Recommended by Sever Dragomir

We give some generalizations of Shafer-Fink inequalities, and prove these inequalities by using a basic differential method and l'Hospital's rule for monotonicity.

Copyright © 2009 W. Pan and L. Zhu. This is an open access article distributed under the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

1. Introduction

Shafer (see Mitrinovic and Vasic [1, page 247]) gives us a result as follows.

Theorem 1.1. *Let $x > 0$. Then*

$$\arcsin x > \frac{6(\sqrt{1+x} - \sqrt{1-x})}{4 + \sqrt{1+x} + \sqrt{1-x}} > \frac{3x}{2 + \sqrt{1-x^2}}. \quad (1.1)$$

The theorem is generalized by Fink [2] as follows.

Theorem 1.2. *Let $0 \leq x \leq 1$. Then*

$$\frac{3x}{2 + \sqrt{1-x^2}} \leq \arcsin x \leq \frac{\pi x}{2 + \sqrt{1-x^2}}. \quad (1.2)$$

Furthermore, 3 and π are the best constants in (1.2).

In [3], Zhu presents an upper bound for $\arcsin x$ and proves the following result.

Theorem 1.3. Let $0 \leq x \leq 1$. Then

$$\begin{aligned} \frac{3x}{2 + \sqrt{1-x^2}} &\leq \frac{6(\sqrt{1+x} - \sqrt{1-x})}{4 + \sqrt{1+x} + \sqrt{1-x}} \leq \arcsin x \\ &\leq \frac{\pi(\sqrt{2} + 1/2)(\sqrt{1+x} - \sqrt{1-x})}{4 + \sqrt{1+x} + \sqrt{1-x}} \leq \frac{\pi x}{2 + \sqrt{1-x^2}}. \end{aligned} \quad (1.3)$$

Furthermore, 3 and π , 6 and $\pi(\sqrt{2} + 1/2)$ are the best constants in (1.3).

Malesevic [4–6] obtains the following inequality by using λ -method and computer separately.

Theorem 1.4. Let $0 \leq x \leq 1$. Then

$$\arcsin x \leq \frac{(\pi(2 - \sqrt{2})) / (\pi - 2\sqrt{2})(\sqrt{1+x} - \sqrt{1-x})}{(\sqrt{2}(4 - \pi)) / (\pi - 2\sqrt{2}) + \sqrt{1+x} + \sqrt{1-x}} \leq \frac{\pi / (\pi - 2)x}{(2 / (\pi - 2)) + \sqrt{1-x^2}}. \quad (1.4)$$

Zhu [7, 8] offers some new simple proofs of inequality (1.4) by L'Hospital's rule for monotonicity.

In this paper, we give some generalizations of these above results and obtain two new Shafer-Fink type double inequalities as follows.

Theorem 1.5. Let $0 \leq x \leq 1$, and $a, b_1, b_2 > 0$. If

$$\begin{aligned} (a, b_1, b_2) \in &\left\{ a \geq 3, b_1 \geq a - 1, b_2 \leq \frac{2a}{\pi} \right\} \\ &\cup \left\{ 3 > a > \frac{\pi}{\pi - 2}, b_2 \leq \frac{2a}{\pi}, b_1 \geq \frac{a \sin t_a}{t_a} - \cos t_a \right\} \\ &\cup \left\{ \frac{\pi}{\pi - 2} \geq a > \frac{\pi^2}{4}, b_2 \leq a - 1, b_1 \geq \frac{a \sin t_a}{t_a} - \cos t_a \right\} \\ &\cup \left\{ \frac{\pi^2}{4} \geq a > 1, b_1 \geq \frac{2a}{\pi}, b_2 \leq a - 1 \right\}, \end{aligned} \quad (1.5)$$

then

$$\frac{ax}{b_1 + \sqrt{1-x^2}} \leq \arcsin x \leq \frac{ax}{b_2 + \sqrt{1-x^2}} \quad (1.6)$$

holds, where t_a is a point in $(0, \pi/2]$ and satisfies $a(t_a \cos t_a - \sin t_a) + t_a^2 \sin t_a = 0$.

Theorem 1.6. Let $0 \leq x \leq 1$, and $c, d_1, d_2 > 0$. If

$$\begin{aligned} (c, d_1, d_2) \in & \left\{ c \geq 6, d_1 \geq c - 2, d_2 \leq \sqrt{2} \left(\frac{2c}{\pi} - 1 \right) \right\} \\ & \cup \left\{ 6 > c > \frac{\pi(2 - \sqrt{2})}{\pi - 2\sqrt{2}}, d_2 \leq \sqrt{2} \left(\frac{2c}{\pi} - 1 \right), d_1 \geq \frac{c \sin t_c}{t_c} - 2 \cos t_c \right\} \\ & \cup \left\{ \frac{\pi(2 - \sqrt{2})}{\pi - 2\sqrt{2}} \geq c > \frac{\pi^2}{8 - 2\pi}, d_2 \leq c - 2, d_1 \geq \frac{c \sin t_c}{t_c} - 2 \cos t_c \right\} \\ & \cup \left\{ \frac{\pi^2}{8 - 2\pi} \geq c > 2, d_1 \geq \frac{\sqrt{2}}{2} \left(\frac{4c}{\pi} - 2 \right), d_2 \leq c - 2 \right\}, \end{aligned} \quad (1.7)$$

then

$$\frac{c(\sqrt{1+x} - \sqrt{1-x})}{d_1 + \sqrt{1+x} + \sqrt{1-x}} \leq \arcsin x \leq \frac{c(\sqrt{1+x} - \sqrt{1-x})}{d_2 + \sqrt{1+x} + \sqrt{1-x}} \quad (1.8)$$

holds, where t_c is a point in $(0, \pi/4]$ and satisfies $c(t_c \cos t_c - \sin t_c) + 2t_c^2 \sin t_c = 0$.

2. One Lemma: L'Hospital's Rule for Monotonicity

Lemma 2.1 (see [9–15]). Let $f, g : [a, b] \rightarrow \mathbb{R}$ be two continuous functions which are differentiable and $g' \neq 0$ on (a, b) . If f'/g' is increasing (or decreasing) on (a, b) , then the functions $(f(x) - f(b))/(g(x) - g(b))$ and $(f(x) - f(a))/(g(x) - g(a))$ are also increasing (or decreasing) on (a, b) .

3. Proofs of Theorems 1.5 and 1.6

(A) We first process the proof of Theorem 1.5.

Let $x = \sin t$ for $x \in (0, 1]$, in which case the proof of Theorem 1.5 can be completed when proving that the double inequality

$$\frac{b_1}{a} \geq \frac{\sin t}{t} - \frac{\cos t}{a} \geq \frac{b_2}{a} \quad (3.1)$$

holds for $t \in (0, \pi/2]$.

Let $F(t) = (\sin t/t) - (\cos t/a)$, we have

$$F'(t) = \frac{t \cos t - \sin t}{t^2} + \frac{\sin t}{a} = \sin t \left(\frac{t \cos t - \sin t}{t^2 \sin t} + \frac{1}{a} \right) =: \sin t \left[H(t) + \frac{1}{a} \right], \quad (3.2)$$

where $H(t) = (t \cos t - \sin t)/(t^2 \sin t) =: f_1(t)/g_1(t)$ and $f_1(t) = t \cos t - \sin t$, $g_1(t) = t^2 \sin t$, $f_1(0) = 0$, $g_1(0) = 0$.

Since $f'_1(t)/g'_1(t) = (-t \sin t)/(2t \sin t + t^2 \cos t) = -(1/(2 + (t/\tan t)))$ decreases on $(0, \pi/2]$, we obtain that $H(t)$ decreases on $(0, \pi/2]$ by using Lemma 2.1. At the same time, $H(0+0) = -1/3$, $H(\pi/2) = -4/\pi^2$, and $F(0+0) = 1 - (1/a)$, $F(\pi/2) = 2/\pi$.

There are four cases to consider.

Case 1 ($a \geq 3$)

Since $F'(t) \leq 0$, $F(t)$ decreases on $(0, \pi/2]$, and $\inf_{x \in (0, \pi/2]} F(t) = 2/\pi$, $\sup_{x \in (0, \pi/2]} F(t) = 1 - 1/a$. So when $b_1 \geq a - 1$ and $b_2 \leq 2a/\pi$, (3.1) and (1.6) hold.

Case 2 ($3 > a > \pi/(\pi - 2)$)

At this moment, there exists a number $t_a \in (0, \pi/2]$ such that $a(t_a \cos t_a - \sin t_a) + t_a^2 \sin t_a = 0$, $F'(t)$ is positive on $(0, t_a]$ and negative on $(t_a, \pi/2]$. That is, $F(t)$ firstly increases on $(0, t_a]$ then decreases on $(t_a, \pi/2]$, and $\inf_{x \in (0, \pi/2]} F(t) = 2/\pi$, $\sup_{x \in (0, \pi/2]} F(t) = F(t_a)$. So when $b_2 \leq 2a/\pi$ and $b_1 \geq a \sin t_a/t_a - \cos t_a$, (3.1) and (1.6) hold.

Case 3 ($\pi/(\pi - 2) \geq a > \pi^2/4$)

Now, $F(t)$ also firstly increases on $(0, t_a]$ then decreases on $(t_a, 2/\pi]$, and $\inf_{x \in (0, \pi/2]} F(t) = 1 - 1/a$, $\sup_{x \in (0, \pi/2]} F(t) = F(t_a)$. So when $b_2 \leq a - 1$ and $b_1 \geq a \sin t_a/t_a - \cos t_a$, (3.1) and (1.6) hold too.

Case 4 ($\pi^2/4 \geq a > 1$)

Since $F'(t) \geq 0$, $F(t)$ increases on $(0, \pi/2]$, $\inf_{x \in (0, \pi/2]} F(t) = 1 - 1/a$, and $\sup_{x \in (0, \pi/2]} F(t) = 2/\pi$. So when $b_1 \geq 2a/\pi$ and $b_2 \leq a - 1$, (3.1) and (1.6) hold.

(B) Now we consider proving Theorem 1.6.

In view of the fact that (1.8) holds for $x = 0$, we suppose that $0 < x \leq 1$ in the following.

First, let $\sqrt{1+x} = \sqrt{2} \cos \alpha$ and $\sqrt{1-x} = \sqrt{2} \sin \alpha$ for $x \in (0, 1]$, we have $x = \cos 2\alpha$ and $\alpha \in [0, \pi/4]$. Second, let $\alpha + \pi/4 = \pi/2 - t$, then $t \in (0, \pi/4]$ and (1.8) is equivalent to

$$\frac{d_1}{c} \geq \frac{\sin t}{t} - \frac{2 \cos t}{c} \geq \frac{d_2}{c}. \quad (3.3)$$

When letting $c = 2a$ and $d_i = 2b_i$ ($i = 1, 2$), (3.3) becomes (3.1).

Let $F(t) = \sin t/t - \cos t/a$. At this moment, $H(t)$ decreases on $(0, \pi/4]$, $H(0+0) = -1/3$, $H(\pi/4) = -(1 - \pi/4)16/\pi^2$, and $F(0+0) = 1 - 2/c$, $F(\pi/4) = \sqrt{2}(2/\pi - 1/c)$.

There are four cases to consider too.

Case 1 ($c \geq 6$)

Since $F'(t) \leq 0$, $F(t)$ decreases on $(0, \pi/4]$, and $\inf_{x \in (0, \pi/4]} F(t) = \sqrt{2}(2/\pi - 1/c)$, $\sup_{x \in (0, \pi/4]} F(t) = 1 - 2/c$. If $d_1 \geq c - 2$ and $d_2 \leq \sqrt{2}(2c/\pi - 1)$, then (3.1) holds on $(0, \pi/4]$ and (1.8) holds.

Case 2 $(6 > c > (\pi(2 - \sqrt{2})) / (\pi - 2\sqrt{2}))$

At this moment, there exists a number $t_a \in (0, \pi/4]$ such that $a(t_c \cos t_c - \sin t_c) + 2t_c^2 \sin t_c = 0$, $F'(t)$ is positive on $(0, t_c]$ and negative on $(t_c, \pi/4]$. That is, $F(t)$ firstly increases on $(0, t_c]$ then decreases on $(t_c, \pi/4]$, and $\inf_{x \in (0, \pi/4]} F(t) = \sqrt{2}((2/\pi) - (1/c))$, $\sup_{x \in (0, \pi/4]} F(t) = F(t_c)$. If $d_2 \leq \sqrt{2}((2c/\pi - 1))$ and $d_1 \geq (c \sin t_c / t_c) - 2 \cos t_c$, then (3.1) holds on $(0, \pi/4]$ and (1.8) holds.

Case 3 $((\pi(2 - \sqrt{2})) / (\pi - 2\sqrt{2})) \geq c > \pi^2 / (8 - 2\pi)$

Now, $F(t)$ also firstly increases on $(0, t_c]$ then decreases on $(t_c, \pi/4]$, and $\inf_{x \in (0, \pi/4]} F(t) = 1 - 2/c$, $\sup_{x \in (0, \pi/4]} F(t) = F(t_c)$. If $d_2 \leq c - 2$ and $d_1 \geq (c \sin t_c / t_c) - 2 \cos t_c$, then (3.1) holds on $(0, \pi/4]$ and (1.8) holds too.

Case 4 $(\pi^2 / (8 - 2\pi)) \geq c > 2)$

Since $F'(t) \geq 0$, $F(t)$ increases on $(0, \pi/4]$, $\inf_{x \in (0, \pi/4]} F(t) = 1 - 2/c$, and $\sup_{x \in (0, \pi/4]} F(t) = \sqrt{2}(2/\pi - 1/c)$. If $d_1 \geq \sqrt{2}(2c/\pi - 1)$ and $d_2 \leq c - 2$, then (3.1) holds on $(0, \pi/4]$ and (1.8) holds.

4. The Special Cases of Theorems 1.5 and 1.6

- (1) Taking $a = 3$, $b_1 = a - 1 = 2$ in Theorem 1.5 and $c = 6$, $d_1 = c - 2 = 4$ in Theorem 1.6 leads to the inequality (1.1).
- (2) Taking $a = \pi / (\pi - 2)$, $b_2 = a - 1 = 2 / (\pi - 2)$ in Theorem 1.5 and $c = (\pi(2 - \sqrt{2})) / (\pi - 2\sqrt{2})$, $d_2 = c - 2 = \sqrt{2}(4 - \pi) / (\pi - 2\sqrt{2})$ in Theorem 1.6 leads to the inequality (1.4).
- (3) Let $a = \pi^2 / 4$, $b_1 = (2/\pi)a = \pi/2$ in Theorem 1.5 and $c = \pi^2 / (2(4 - \pi))$, $d_1 = (2\sqrt{2}/\pi)c - \sqrt{2} = 2\sqrt{2}(\pi - 2) / (4 - \pi)$ in Theorem 1.6, we have the following result.

Theorem 4.1. Let $0 \leq x \leq 1$. Then

$$\frac{(\pi^2/4)x}{\pi/2 + \sqrt{1-x^2}} \leq \frac{(\pi^2/(8-2\pi))(\sqrt{1+x} - \sqrt{1-x})}{2\sqrt{2}(\pi-2)/(4-\pi) + \sqrt{1+x} + \sqrt{1-x}} \leq \arcsin x. \quad (4.1)$$

Furthermore, $\pi^2/4$ and $\pi/2$, $\pi^2/(8-2\pi)$ and $2\sqrt{2}(\pi-2)/(4-\pi)$ are the best constants in (4.1).

References

- [1] D. S. Mitrinović and P. M. Vasic, *Analytic Inequalities*, vol. 16 of *Grundlehren der mathematischen Wissenschaften*, Springer, New York, NY, USA, 1970.
- [2] A. M. Fink, "Two Inequalities," *Publikacije Elektrotehničkog Fakulteta Univerziteta u Beogradu. Serija Matematika*, vol. 6, pp. 48–49, 1995.
- [3] L. Zhu, "On Shafer-Fink inequalities," *Mathematical Inequalities & Applications*, vol. 8, no. 4, pp. 571–574, 2005.
- [4] B. J. Malešević, "One method for proving inequalities by computer," *Journal of Inequalities and Applications*, vol. 2007, Article ID 78691, 8 pages, 2007.

- [5] B. J. Malešević, "An application of λ -method on inequalities of Shafer-Fink's type," *Mathematical Inequalities & Applications*, vol. 10, no. 3, pp. 529–534, 2007.
- [6] B. J. Malešević, "Some improvements of one method for proving inequalities by computer," preprint, 2007, <http://arxiv.org/abs/math/0701020>.
- [7] L. Zhu, "On Shafer-Fink-type inequality," *Journal of Inequalities and Applications*, vol. 2007, Article ID 67430, 4 pages, 2007.
- [8] L. Zhu, "New inequalities of Shafer-Fink type for arc hyperbolic sine," *Journal of Inequalities and Applications*, vol. 2008, Article ID 368275, 5 pages, 2008.
- [9] G. D. Anderson, M. K. Vamanamurthy, and M. Vuorinen, "Inequalities for quasiconformal mappings in space," *Pacific Journal of Mathematics*, vol. 160, no. 1, pp. 1–18, 1993.
- [10] G. D. Anderson, M. K. Vamanamurthy, and M. K. Vuorinen, *Conformal Invariants, Inequalities, and Quasiconformal Maps*, Canadian Mathematical Society Series of Monographs and Advanced Texts, John Wiley & Sons, New York, NY, USA, 1997.
- [11] G. D. Anderson, S.-L. Qiu, M. K. Vamanamurthy, and M. Vuorinen, "Generalized elliptic integrals and modular equations," *Pacific Journal of Mathematics*, vol. 192, no. 1, pp. 1–37, 2000.
- [12] I. Pinelis, "L'Hospital type results for monotonicity, with applications," *Journal of Inequalities in Pure and Applied Mathematics*, vol. 3, no. 1, article 5, pp. 1–5, 2002.
- [13] L. Zhu, "Sharpening Jordan's inequality and the Yang Le inequality," *Applied Mathematics Letters*, vol. 19, no. 3, pp. 240–243, 2006.
- [14] D.-W. Niu, Z.-H. Huo, J. Cao, and F. Qi, "A general refinement of Jordan's inequality and a refinement of L. Yang's inequality," *Integral Transforms and Special Functions*, vol. 19, no. 3–4, pp. 157–164, 2008.
- [15] S. Wu and L. Debnath, "A generalization of L'Hôpital-type rules for monotonicity and its application," *Applied Mathematics Letters*, vol. 22, no. 2, pp. 284–290, 2009.